

# The Challenges in Spectral Image Analysis: an Introduction, and Review of ANN Approaches \*

Erzsébet Merényi

University of Arizona, Lunar and Planetary Laboratory, PIRL  
Tucson, AZ 85721, U.S.A.

**Abstract.** Utilization of remote sensing multi- and hyperspectral imagery has been rapidly increasing in numerous areas of economic and scientific significance. Hyperspectral sensors, in particular, provide the detailed information that is known from laboratory measurements to characterize and identify minerals, soils, rocks, plants, water bodies, and other surface materials. This opens up tremendous possibilities for resource exploration and management, environmental monitoring, natural hazard prediction, and more. However, exploitation of the wealth of information in spectral images has yet to match up to the sensors' capabilities, as conventional methods often prove inadequate. ANNs hold the promise to revolutionize this area by overcoming many of the mathematical obstacles that traditional techniques fail at. By providing high speed when implemented in parallel hardware, (near-)real time processing of extremely high data volumes, typical in remote sensing spectral imaging, will also be possible.


## 1. Challenges in remote spectral image analyses

Airborne and satellite-borne spectral imaging has become one of the most advanced tools for collecting vital information about the surface covers of Earth and other planets. The utilization of these data includes areas such as mineral exploration, land use, forestry, natural hazard assessments, water resources, environmental contamination, ecosystem management, biomass and productivity assessment, and many other activities of economic significance, as well as prime scientific pursuits such as looking for possible sources of past or present life on other planets. The number of applications has dramatically increased in the past ten years with the advent of imaging spectrometers, which greatly surpass traditional multi-spectral imagers (*e.g.*, Landsat Thematic Mapper) in that they can resolve the detailed spectral features that are known to characterize minerals, soils, rocks, and vegetation, from laboratory measurements. While a multi-spectral sensor samples the given wavelength window (typically the 0.4 – 2.5

---

\*Work funded by NASA, Applied Information Systems Research Program, NAG54001; Paper in Proceedings of the 7th European Symposium on Artificial Neural Networks, ESANN99, Bruges, Belgium, April 21–23, 1999, pp 93–98.

$\mu\text{m}$  range in the case of Visible and Near-Infrared surface reflectance imaging) with several broad bandpasses, leaving large gaps between the bands, spectral imagers (hyperspectral sensors) sample the spectral window contiguously with very narrow bandpasses. Figures 1 and 2 illustrate the above.



**Figure 1 (left).** The concept of spectral imaging. Figure from Campbell (1996) [1]. **Figure 2 (right).** The spectral signature of the mineral alunite as seen through the 6 broad channels of Landsat TM, as seen by the moderate spectral resolution sensor MODIS, and as measured in the laboratory. Hyperspectral sensors such as AVIRIS of NASA/JPL [2] provide spectral details comparable to laboratory measurements.

Formally, the vector  $S^{x,y} = (S_1^{x,y}, \dots, S_{NB}^{x,y})$ , where  $S_k^{x,y}$  is the data value in the  $k$ th image band ( $k = 1, \dots, NB$ ) at pixel location  $(x, y)$ , is called a spectrum. It is a characteristic pattern (Figures 1, 2) which provides a clue to the surface material(s) within pixel  $(x, y)$ .  $NB$  denotes the number of image bands. The feature space spanned by VIS-NIR reflectance spectra is  $[0, U]^{NB} \subset \mathfrak{R}^{NB}$  where  $U > 0$  represents an upper limit of the measured scaled reflectivity. Sections of this space can be very densely populated while other parts may be extremely sparse, depending on the materials in the scene and on the spectral resolution of the sensor. Great spectral detail comes at a cost of a very high data volume, and it also poses new mathematical challenges in the classification of images with high spectral dimensionality. The specific problems associated with remote sensing spectral image analyses arise from any combination of the following [3]:

- The spectral patterns are high dimensional (dozens  $\leq NB \leq$  hundreds);
- The number of data points (image pixels) can be as large as several millions;
- The pixels are mixed: Several different materials contribute to the spectral signature detected from each pixel;
- Given the richness of data, the goal is to separate many cover classes;
- Different surface materials may be distinguished by very subtle differences in their spectral patterns;
- Very little training data may be available for some classes; and classes may be represented very unevenly.

ANNs have been gaining recognition as powerful answers to the above challenges. It should be emphasized that traditional lower dimensional multi-spectral images also benefit greatly from ANN algorithms because remote sensing spectral images of any dimensionality share all but the first problem above.

For this discussion, we will omit additional effects such as atmospheric distortions, illumination geometry and albedo variations in the scene, because these can be addressed through well-established procedures prior to classification.

## 2. A review of ANN approaches and results

In the following, emphasis is on works that overcome mathematical obstacles, or improve classification quality over conventional algorithms. In particular, the speed benefit of parallel hardware implementations is not discussed. For sake of space, the reader is referred to the bibliographies in the referenced papers for relevant further works, including well-known ANN paradigms.

Traditional multi-spectral images (*e.g.*, Landsat TM) have long been shown to gain improved accuracy from ANN classifications, using BP networks [4-6], or variants of self-organization, vector quantization, and their hybrids [7-8]. ART networks and variants by [9] were successful in distinguishing vegetation species.

Much less work has been done with hyperspectral images, although this type of data would clearly be much better exploited with ANNs than with classical methods. The reason for little work in this area is a combination of the novelty of hyperspectral imaging (< 10 years in comparison to over 25 years of Landsat), and that the high dimensionality of the input data space requires large, complex networks. [10] presented one of the pioneering papers on simulated 201-band spectra, which were reduced to 20, 40 and 60 bands using feature extraction prior to classification into 3 classes. Comparison of several classifiers including Maximum Likelihood (ML), BP network and a Parallel Self-organizing Hierarchical Neural Network (PSHNN) favored the ML, with PSHNN next. However, the authors admitted that the ML had an advantage by virtue of gaussian data generation. [11] successfully classified a real AVIRIS image of the Neovolcanic Zone in South-Central Iceland into 9 geological classes, reducing first the 224 AVIRIS bands to 35. As an important advantage over traditional feature extractors such as PCA, they used an ANN (the same network that performed the classification itself) for Decision Boundary Feature Extraction (DBFE). The DBFE is claimed to preserve all features that are necessary to achieve the same accuracy as in the original data space, by a given classifier.

Self-Organizing Maps have been recognized as useful tools for classification of images with high spectral dimension. For supervised case, the general observation is that an SOM component in the ANN architecture makes network training much easier (than, *e.g.*, training a BP network); that it produces more accurate classification results based on a smaller amount of training spectra than would be required for the training of BP [10]. Using a hybrid SOM-BP architecture, [12] mapped previously undetected soil variants on Mars from 90-band images, [13] improved asteroid compositional taxonomy from 65-band spectra, with no

prior feature extraction. Full spectral resolution AVIRIS images were classified into large number of output classes by a similar approach [3].

For discoveries in data spaces, SOMs have also been successfully used for the detection of surface compositional classes that were missed by PCA or other conventional techniques [13–14]. MacDonald *et al.*, [15] compare three unsupervised techniques: the Kohonen SOM, the Scale Invariant Feature Map, and the Generative Topographic Mapping, which is a “principled alternative to the SOM”. They arrive at similar preliminary results as [13], on the same 65-D data. Since convergence of the GTM can be proven and it has a well-defined cost function, this investigation may develop into better understanding of hyperspectral spaces than was gained by the above previous works.

The mixed pixel problem is addressed by Pendock [16], using an associative ANN to establish a linear mixture model for the areal contributions of “endmember” materials in each pixel. (The endmembers are the spectra whose weighted sum makes up the spectral signature of each pixel. These are typically not the same as the Principle Components of the spectral image.) The linear unmixing approach is one of the most popular conventional techniques in interpreting spectral images [17]. Automated determination of the endmembers, however, has not been very successful. Pendock’s approach [16] brings a new solution.

In remote sensing, obtaining an ideal number of reliable training samples can be hindered by the inaccessibility of certain locations, or by the fact that small outcrops of important materials may contain very few recognizable pixels in the scene [3]. This can render some of the most valued conventional classifiers, notably covariance based ones such as ML, useless because those require at least  $NB + 1$  samples for each class [10], [3]. Identification of  $NC * (NB + 1)$  samples, where  $NC$  is the number of surface cover classes, can be prohibitively expensive, or impossible, for large  $NB$  and  $NC$ . Fardanesh and Ersoy [18] offer an architectural approach to compensate for small training sets.

Many believe that hyperspectral images are highly redundant because of band correlations. Others maintain an opposite view. Few investigations exist yet into intrinsic dimensionality (ID) of hyperspectral images. Bruske [19] finds the spectral ID of an AVIRIS image to be between 4 and 7, using Optimally Topology Preserving Maps. This seems consistent with the number of mixture model endmembers in many works, and can be a step toward understanding spectral image compression. The number of separable meaningful spectral classes is an important related question, the answer to which was seen to be more complex when ANNs were utilized than with conventional methods [13–14]. [20] presents a Growing SOM approach applied to Landsat imagery, which can provide more theoretical insight as well as a better practical handle on cluster determination.

Visualisation of data clusters in higher-dimensional spaces as detected by SOM type mappings has been targeted by several works [21–24], however, no application to hyperspectral images has been published yet. Merényi engineered a tool specific to hyperspectral data, utilizing [21–22] and [24], and detection of  $\sim 30$  geologically meaningful clusters in an AVIRIS 194-band image from its

SOM is demonstrated at <http://www.arizona.edu/~erzsebet/annps.html> .

### 3. Future work, outstanding problems

Application of ANNs to spectral, especially to hyperspectral imagery is in its infancy. Further, robust solutions are urgently needed to the above, as well as to some other closely related issues (not discussed here) such as classification of multi-source disparate data in conjunction with spectral images [9]; missing bands; variable spectral resolution.

Image compression has an even more pressing significance for multi- and hyperspectral data than for monochrome and RGB images. Transmission of enormous data volumes from satellites with limited downlink capacity, and storage of these data merit serious considerations. Encouraging improvement over the most widely used JPEG compression algorithm is presented by Amerijckx *et al.*, [25] using an SOM. Hopefully, such approaches can be extended to multispectral imagery in a way that takes into account and makes use of the band correlations.

### References

1. J.Campbell: *Introduction to Remote Sensing*. The Guilford Press (1996)
2. R.O. Green, Ed.: *Summaries of 6th Annual JPL Airborne Geoscience Workshop, Vol. 1. AVIRIS Workshop*. Pasadena, CA, March 4–6, 1996 (1996)
3. E. Merényi: Self-Organizing ANNs for Planetary Surface Composition Research. *Proc. 6th European Symposium on Artificial Neural Networks, ESANN'98, Bruges, Belgium, April 22–24, 1998*, 197–202 (1998)
4. G.F. Hepner, T. Logan, N. Ritter, N. Bryant: Artificial Neural Network Classification Using a Minimal Training Set: Comparison to Conventional Supervised Classification. *Photogrammetric Eng. & Remote Sensing*, 56(4), 469–473 (1990)
5. J.D. Paola, R.A. Schowengerdt: Comparison of Neural Network to Standard Techniques for Image Classification and Correlation. *Proc. Int'l Geosci. and Remote Sensing Symposium, Caltech, Pasadena, CA, August 8–12, 1994*, III, 1404–1405, (1994)
6. Y. Wang, Y. and D.L. Civco: Artificial Neural Networks in High Dimensional Spatial Data Classification: A Performance Evaluation. *ACMS/ASPRS Annual Convention & Exposition Technical Papers*, 3, 662–671, Charlotte North Carolina, February 27 – March 2 1995 (1995)
7. E. Merényi, B. Csathó, M. Bodrogi, Á. Gulyás: Utilization of Landsat Images For Mapping Natural Resources and For Environmental Protection In Hungary. *Proc. Tenth Thematic Conference on Geologic Remote Sensing, San Antonio, TX, 9–12 May, II*, 491–502 (1994)
8. B. Solaiman and M.C. Mouchot: A Comparative Study of Conventional and Neural Network Classification of Multispectral Data. *Proc. Int'l Geosci. and Remote Sensing Symposium, Caltech, Pasadena, CA, August 8–12, 1994*, III, 1413–1415, (1994)

9. G.A. Carpenter, M.N. Gajja, S. Gopal, and C.E. Woodcock: ART Neural Networks for Remote Sensing:Vegetation Classification from Landsat TM and Terrain Data. *IEEE. Trans. Geosci. and Remote Sens.*, 35(2), 308–325 (1997)
10. J.A. Benediktsson, P.H. Swain, O.K. Ersoy, D. Hong: Classification of Very High Dimensional Data Using Neural Networks. *IGARSS'90 10th Annual International Geoscience and Remote Sensing Symp.*, 2, 1269 (1990)
11. J.A. Benediktsson, J.R. Sveinsson, K. Arnason, K.: Classification of Very-High-Dimensional data With Geological Applications. *Proc. MAC Europe 91, Lenggreis, Germany, 4–6 October, 1994*, 13–18 (1994)
12. E. Merényi, R.B. Singer, J.S. Miller: Mapping of Spectral Variations On the Surface of Mars From High Spectral Resolution Telescopic Images. *ICARUS*, 124, 280–295 (1996a)
13. E.S. Howell, E. Merényi, L.A. Lebofsky: Classification of Asteroid Spectra Using a Neural Network. *J. Geophys. Res.*, 99, 10,847–10,865 (1994)
14. E. Merényi, E.S. Howell, L.A. Lebofsky, A.S. Rivkin: Prediction of Water In Asteroids From Spectral Data Shortward of 3 Microns. *ICARUS*, 129, 421–439 (1997a)
15. D. MacDonald, S. McGlinchey, J. Kawala, and C. Fyfe: Comparison of Kohonen, Scale-Invariant and GTM Self-organizing Maps for Interpretation of Spectral Data. *This volume.*, (1999)
16. N. Pendock: A Simple Associative Neural Network for Producing Spatially Homogeneous Spectral Abundance Interpretations of Hyperspectral Imagery. *This volume.*, (1999)
17. B. Adams, M.O. Smith, and A.R. Gillespie (1993), “Imaging spectroscopy: Interpretation based on spectral mixture analysis,” In *Remote Geochemical Analysis: Elemental and Mineralogical Composition*, C.M. Peters and P.A.J. Englert, Eds., Cambridge University Press, New York. pp. 145–166.
18. M.T. Fardanesh and O.K. Ersoy: Classification Accuracy Improvement of Neural Network Classifiers by Using Unlabeled Data. *IEEE Trans. Geosci. and Remote Sens.*, 36(3), 1020–1025 (1998)
19. J. Bruske, and E. Merényi: Estimating the Intrinsic Dimensionality of Hyperspectral Images. *This volume.*, (1999)
20. T. Villmann: Benefits and Limits of the Self-Organizing Map and its Variants in the Area of Satellite Remote Sensing Processing. *This volume.*, (1999)
21. A. Ultsch, and H. P. Simeon: Kohonen’s Self Organizing Feature Map for Exploratory Data Analysis. *Proc. INNC-90-PARIS*, I, 305–308 (1990)
22. M. Cottrell, E. de Bodt: A Kohonen Map Representation to Avoid Misleading Interpretations. *European Symposium on Artificial Neural Networks, Bruges, Belgium, 22–24 April, 1996*, (1996)
23. M.A. Kraaijveld, J. Mao, and A.K. Jain: A Nonlinear Projection Method Based on Kohonen’s Topology Preserving Maps. *IEEE Trans. on Neural Networks*, 6(3), 548–559 (1995)
24. S. Kaski, J. Nikkila, and T. Kohonen: Methods for Interpreting a Self-Organizing Map in Data Analysis. *Proc. 6th European Symposium on Artificial Neural Networks, ESANN'98, Bruges, Belgium, April 22–24, 1998*, 185–190 (1998)
25. C. Amerijckx, M. Verleysen, P. Thissen, and J-D. Legat: Image Compression by Self-Organized Kohonen Map. *IEEE Trans. on Neural Networks*, 9(3), 503–507 (1998)